


Agroinfiltration basics

Animal and plant cell culture
Practicum: Plant tissue culture
2024/2025


Virulent strains of *Agrobacterium tumefaciens*

T-DNA transfer is mediated by products encoded by the 30-40 kb *vir* region of the Ti plasmid (a large plasmid, more than 200 kb, with a key role in tumor induction). This region consists of at least six essential operons (*vir A*, *vir B*, *vir C*, *vir D*, *vir E*, *virG*) and two non-essential ones (*virF*, *virH*). The only constitutively expressed operons are *virA* and *virG*, which encode a two-component system (VirA-VirG) that activates transcription of the other *vir* genes.




The T-DNA contains two types of genes: the oncogenic genes coding for enzymes involved in the synthesis of auxins and cytokinins and responsible for tumor formation, and the genes coding for the synthesis of opines that are produced and secreted by the crown gall cells and consumed by *A. tumefaciens* as a source of carbon and nitrogen.

Induction of the bacterial virulence system



Generation of the T-DNA transfer complex




Activation of the vir genes leads to the generation of single-stranded (ss) molecules that represent the copy of the bottom T-DNA strand. Any DNA located between the **T-DNA borders** will be transferred into the plant cell as single-stranded DNA and integrated into the plant genome.

The proteins VirD1 and VirD2 play a key role in this step. They recognize the T-DNA border sequences and nick (endonuclease activity) the bottom strand at each border.

After endonucleotidic cleavage, VirD2 remains covalently bound to the 5' end of the ss T-DNA strand. This association prevents exonucleolytic attack on the 5'-end of the ss-T-DNA strand and marks the 5'-end as the leading end of the T-DNA transfer complex.


Translocation of the T-DNA complex



The ssT-DNA-protein complex is transferred into the plant cell. According to the most widely accepted model, the ssT-DNA-VirD2 complex is enveloped by the VirE2 protein, a single-stranded DNA-binding protein. This cooperative association prevents attack by nucleases and also elongates the ssT DNA strand, reducing the diameter of the complex to about 2 nm, which facilitates translocation through membrane channels.

VirB are membrane-associated proteins and most VirB proteins are assembled into a membrane-spanning protein channel that spans both membranes.


Integration of the T-DNA into the plant genome



Within the plant cell, the ssT-DNA complex is transported through the nuclear membrane into the nucleus. It was found that two Vir proteins play an important role in this step: *VirD2* and *VirE2*. It is hypothesized that the two NLSs of *VirE2* are important for the continuous nuclear import of the ssT-DNA complex.

The final step of T-DNA transfer is its integration into the plant genome. It is assumed that integration occurs by illegitimate recombination. The cleavage of some bases confers only minimal specificity to the recombination process by positioning *VirD2* for ligation.


Agrobacterium tumefaciens strain GV3101



The *Agrobacterium tumefaciens* strain GV3101 used for transformation harbors the C58 chromosomal backbone containing **rifampicin** resistance and carries two plasmids. A non-oncogenic disarmed tumor-inducing plasmid (Ti plasmid) containing the virulence (*vir*) genes and resistance to **gentamicin**, but lacking the T-DNA region. The T-DNA region is present on a second plasmid, the binary vector. The selection marker in the binary vector is the **kanamycin** resistance gene. DNA sequences to be transformed are cloned between the left and right border sequences of the T-DNA region and transferred to the host.

Binary vectors

A binary vectors are standard tool in the transformation of higher plants mediated by *Agrobacterium tumefaciens*. It is composed of the borders of T-DNA, multiple cloning sites, replication functions for *Escherichia coli* and *A. tumefaciens*, selectable marker genes, reporter genes, and other accessory elements that can improve the efficiency of and/or give further capability to the system.




Binary vectors

A binary vectors are standard tool in the transformation of higher plants mediated by *Agrobacterium tumefaciens*. It is composed of the **borders of T-DNA**, multiple cloning sites, replication functions for *Escherichia coli* and *A. tumefaciens*, selectable marker genes, reporter genes, and other accessory elements that can improve the efficiency of and/or give further capability to the system.




Binary vectors

A binary vectors are standard tool in the transformation of higher plants mediated by *Agrobacterium tumefaciens*. It is composed of the **borders of T-DNA**, multiple cloning sites, **replication functions** for *Escherichia coli* and *A. tumefaciens*, selectable marker genes, reporter genes, and other accessory elements that can improve the efficiency of and/or give further capability to the system.




Binary vectors

A binary vectors are standard tool in the transformation of higher plants mediated by *Agrobacterium tumefaciens*. It is composed of the **borders of T-DNA**, multiple cloning sites, **replication functions** for *Escherichia coli* and *A. tumefaciens*, **selectable marker genes**, reporter genes, and other accessory elements that can improve the efficiency of and/or give further capability to the system.




Binary vectors

A binary vectors are standard tool in the transformation of higher plants mediated by *Agrobacterium tumefaciens*. It is composed of the **borders of T-DNA**, multiple cloning sites, **replication functions** for *Escherichia coli* and *A. tumefaciens*, **selectable marker genes**, **reporter genes**, and other accessory elements that can improve the efficiency of and/or give further capability to the system.




Agrobacterium tumefaciens strain GV3101

Acetosyringone




Agroinfiltration





Nicotiana benthamiana

Agroinfiltration




Nicotiana benthamiana

Agroinfiltration



Nicotiana benthamiana

Confocal microscopy after argoinfiltration with plasmids encoding GFP and RFP-tagged proteins, respectively.



Nicotiana benthamiana
leaf epidermal cells.

Cell viability
assays




```
graph TD; A[Cell viability assays] --> B[FDA]; A --> C[PI];
```

The diagram is a simple flowchart. At the top center is a light gray rounded rectangle containing the text 'Cell viability assays'. Two thick black arrows point downwards from the bottom corners of this rectangle to two separate black rounded rectangles below. The left rectangle contains the text 'FDA' in green, and the right rectangle contains the text 'PI' in red.

FDA


PI

Cell viability assay - fluorescein diacetate (FDA)




FDA enters the cells, which then convert it into fluorescein. FDA is a colorless molecule that does not fluoresce, and fluorescein fluoresces green when illuminated with blue light. Since only living cells have active esterases that can metabolize FDA, the green fluorescence within the cell is used as an **indicator of viable cells**.

Cell viability assay - propidium iodide (PI)




PI is a dye that **intercalates into the DNA of dead cells**, as it cannot pass through the cell membranes of living cells, and fluoresces red in green light. PI is also used in plant biology to make cell walls visible.



Coskun et al. Journal of experimental botany (2011) 63(1), 151–162

Confocal micrographs showing propidium iodide staining of the cell wall and nuclei of damaged cells from lateral root tips of intact barley (*Hordeum vulgare* L.) seedlings treated with silver ions.



Application of propidium iodide for the labeling of cell walls in the analysis of GFP-tagged PIN1 and PIN7 protein localization.